Milwaukee County COVID-19 Data Summary

Milwaukee County COVID-19 Epidemiology Intel Team

This report was updated on October 29, 2020 and includes data through October 27, 2020. Note that data for recent weeks may be under-reported due to pending test results.

Milwaukee County COVID-19 Summary Statistics

Overall Milwaukee County COVID-19 Summary Statistics March 1 - October 27					
	Milwaukee County	City of Milwaukee	Suburbs		
Total tests performed	553,401	359,166	194,235		
Percent positive of all tests performed	8.4%	8.9%	7.5%		
Number of confirmed cases	40,705	28,202	12,503		
Number of hospitalizations	2,952	2,080	872		
Number of deaths	572	356	216		
Case fatality rate	1.4%	1.3%	1.7%		

Weekly Milwaukee County COVID-19 Summary Statistics October 21 - October 27				
	Milwaukee County	City of Milwaukee	Suburbs	
Total tests performed	25,597	15,478	10,119	
Percent positive of all tests performed	14.0%	14.2%	13.8%	
Number of confirmed cases	3,017	1,910	1,107	
Number of hospitalizations	183	99	84	
Number of deaths	7	2	5	

Total Cases and New Cases

There are now a total of 40705 cases in Milwaukee County, since the first confirmed case on March 6th, 2020. Over the last week, we observed 3017 new confirmed cases in Milwaukee County, including 1910 new cases in the city of Milwaukee. **Figure 1** shows the daily incidence of new cases (bars) and the average daily incidence within the last 7 days (line), which provides a smoothing effect to enhance visualization, for both the city and the county. To indicate a potential reporting delay, we shade the last four days of data and exclude those days from the trend line.

Over the last week, we have seen a continued increase in confirmed cases. The highest daily case count since the beginning of the epidemic occurred on October 19, 2020, with 674 cases in the county overall. The highest daily case count over the entire epidemic in the suburbs occurred on October 19, 2020, with a total of 264 cases confirmed. The highest case count in the city occurred on October 23, 2020, with a total of 426 cases confirmed.

Milwaukee County Suburbs

City of Milwaukee

200

100

200

100

And John St. John S

Figure 1: Milwaukee County daily number of COVID-19 cases

Date of specimen collection

Data source: Wisconsin Electronic Disease Surveillance System (WEDSS) Created by the Milwaukee County COVID-19 Epidemiology Intel Team

Total Deaths and New Deaths

There are a total of 572 COVID-19 related deaths in Milwaukee County. Over the last week, we observed 7 deaths, with 2 from the City of Milwaukee. **Figure 2** shows the number of daily COVID-19 related deaths among Milwaukee County and City of Milwaukee residents. The overlaid lines show the average daily deaths within the last 7 days for each jurisdiction. Overall, daily deaths have remained relatively stable since mid-July. Deaths peaked on April 8, 2020. Several smaller peaks in deaths are notable since April 8th, with an overall decrease to a low number of deaths beginning in early June. Over the last week, the number of deaths in the suburbs has increased slightly.




Figure 2: Milwaukee County COVID-19 daily deaths

Data source: Wisconsin Electronic Disease Surveillance System (WEDSS) Created by the Milwaukee County COVID-19 Epidemiology Intel Team

The COVID-19 Growth Rate

The time it takes for the number of cases to double is called the doubling time. **Figure 3** shows doubling times for Milwaukee County, the surrounding Waukesha, Ozaukee and Walworth (WOW) counties, the M7 (7-county) metropolitan area, and the state of Wisconsin. Dotted lines indicate doubling times of 1, 2 3 and 4 days, which are generally associated with a condition of exponential growth. The current doubling time in Milwaukee County is 100.00 days. The current doubling time for WOW counties is 59.67 days. The current doubling time for the state of Wisconsin is 42.25 days. **Figure 4** shows the trend in doubling times for Milwaukee County and the City of Milwaukee as compared to the state, over the course of the epidemic. As illustrated, the epidemic initially doubled more quickly in Milwaukee County and the city, but has since slowed (improved) more in the city and county than in the state as a whole.

Figure 3: Cumulative cases after 20 confirmed Confirmed cases (log10 scale) MKE County 100000 MKE Metro **WOW Counties** 10000 WI State 1-day doubling 1000 2-day doubling 3-day doubling 100 4-day doubling 100 120 140 160 180 200 220 20 40 60 Days since 20 confirmed cases

Data source: Wisconsin Department of Health Services Created by the Milwaukee County COVID-19 Epidemiology Intel Team




Figure 4: Trend in doubling times

Data sources: WI Department of Health Services & WI Electronic Disease Surveillance System Created by the Milwaukee County COVID-19 Epidemiology Intel Team

The COVID-19 Reproductive Number

Another way of examining the growth rate of the infection is to examine the reproductive number (R). This number captures the number of new cases that are the result of an existing case. For example, an R of 2 would indicate that each infected person infects 2 new people. The following plots show the change in R over time for Milwaukee County, **Figure 5**, the City of Milwaukee, **Figure 6a**, and Milwaukee County suburbs, **Figure 6b**. Each plot includes key dates related to physical distancing or focused testing campaigns affecting residents. The R for each date is calculated to represent the R for a 7-day period with the start day of that 7-day period represented on the graph.

After the first minimum R value in Milwaukee County observed (R = 0.84 on April 7, 2020), we observed an increase in R to a high of 1.52 on May 8, 2020 and then a decrease to a low of 0.76 in the county on May 21, 2020. The R increased again to a recent high of 1.41 on September 11, 2020. Patterns in the City of Milwaukee are very similar to those in the county overall. Patterns in the suburbs show more fluctuation. The R values for the most recent week are 1.040 for the county, 1.055 in the city, and 1.016 in the suburbs.


5.0 May 10: Mother's Day Aug 1 Oct 8 4th of July holiday First distancing Easter & Statewide WI state May 11: Two free test sites open 4.5 Passover weekend mask order begins order 25% May 13: WI Supreme Court ruling Reproductive number, capacity 4.0 May 22: Suburbs partially re-open March 25 April 18-20 July 10 Aug 18-Sept 8 . HOC testing Safer Several schools May 25: Memorial Day Shorewood 3.5 nask order begins May 26: City partially re-opens April 24-28 July 16 Sept 7 3.0 Smithfield City of MKE _abor Day estina mask order begins June 5 2.5 WI election City restaurants 2.0 1.5 1.0 0.5 0.0 Jun-01 Jul.O7 AUGO OT mu'nu'nu'; Mr. Mr. Mr. V Section Sectio

Figure 5: One week reproductive number for Milwaukee County

One-week window start date

Data source: Wisconsin Electronic Disease Surveillance System (WEDSS) Created by the Milwaukee County COVID-19 Epidemiology Intel Team


Figure 6a: One week reproductive number for City of Milwaukee



One-week window start date

Data source: Wisconsin Electronic Disease Surveillance System (WEDSS) Created by the Milwaukee County COVID-19 Epidemiology Intel Team

Figure 6b: One week reproductive number for Milwaukee County suburbs



One-week window start date

Data source: Wisconsin Electronic Disease Surveillance System (WEDSS) Created by the Milwaukee County COVID-19 Epidemiology Intel Team


Demographic Patterns – Age, Sex, Race and Ethnicity

Confirmed cases

COVID-19 cases vary by demographic characteristics. **Figure 7** shows cumulative case plots including confirmed positive cases with an available specimen collection date, plotted by census block group (CBG) median household income, sex, age, and race/ethnicity groups. Most diagnosed cases fall within the ages of 18-79. Of all confirmed cases, 46% are male and 53% are female. The largest number of cases have been identified among the non-Hispanic White population (N = 14965), followed by the Hispanic population (N = 11235), and the Black/AA population (N = 9377). The lower two quartiles of median household income (\$0 - \$35,833, and \$35,834 to \$50,096) have a larger number of cases than the higher two quartiles (\$50,097 to \$68,393, and \$68,394 to \$250,001), with the fewest cases identified among the highest income group.

Over the past week, we have observed increases among individuals in all income groups, most age groups, and among non-Hispanic Whites, non-Hispanic Black/AAs, and Hispanics, with similar increases for both sexes. The cumulative number of cases among those ages 25-39 (N = 12351) still exceeds the number among the next highest group, those ages 40-59 (N = 10899). In the last few weeks, we saw the number of cases under age 18 (N = 4465) increase to approach the number diagnosed among those 60-79 (N = 4924).

Figure 7: Cumulative confirmed cases in Milwaukee County



Date of specimen collection


Data source: Wisconsin Electronic Disease Surveillance System (WEDSS) Created by the Milwaukee County COVID-19 Epidemiology Intel Team

*Race and ethnicity were combined into one variable where the Hispanic category includes Hispanics of any race.

AIAN stands for American Indian or Alaska Native and NHOPI stands for Native Hawaiian or Other Pacific Islander.


When examined as population-based rates in **Figure 8**, demographic patterns are also apparent. Early in the epidemic, we saw a clear age gradient in population-based rates, with older populations experiencing greater rates. However, in the last months, we have seen rates among the younger, working age groups (18-24, 25-39, and 40-59) exceed the rate of those aged 80+. The rate among those 60-79 now approaches that of those aged 80+ and the rate among those 40-59 exceeds that among those 18-24. By race and ethnicity, the rate was highest among Black/AA populations until the beginning of May, when we observed a surge among Hispanics resulting in the Hispanic rate (79.92 per 1,000 people) exceeding that among all other racial and ethnic groups. The rate among AIANs (40.89 per 1,000 people) is now slightly higher than that among Black/AAs (37.66 per 1,000 people), although the total number of cases among AIANs (N = 190) is much lower than the number among Black/AAs (N = 9377). The rate among Asians comes next, at 33.92 per 1,000 people. The population-based rate among non-Hispanic Whites is relatively low compared to other racial and ethnic groups. Rates are very similar among males and females.

Figure 8: Population based incidence rates in Milwaukee County




Age	N Cases	Population	Rate per 1,000 residents
< 18	4465	231111	19.32
18-24	6800	150895	45.06
25-39	12351	163246	75.66
40-59	10899	230887	47.20
60-79	4924	142783	34.49
80+	1266	35287	35.88

Specimen collection date



Race/Ethnicity*	N Cases	Population	Rate per 1,000 residents
Black or AA	9377	249011	37.66
White	14965	493723	30.31
Hispanic	11235	140575	79.92
Asian	1372	40443	33.92
AIAN	190	4647	40.89
Multiple Races	165	24224	6.81

Specimen collection date



Gender	N Cases	Population	Rate per 1,000 residents
Male	18882	461670	40.90
Female	21728	492539	44.11

Specimen collection date

Data source: Wisconsin Electronic Disease Surveillance System (WEDSS)

Created by the Milwaukee County COVID-19 Epidemiology Intel Team

*Race and ethnicity were combined into one variable where the Hispanic category includes Hispanics of any race.

AIAN stands for American Indian or Alaska Native and NHOPI stands for Native Hawaiian or Other Pacific Islander.

Hospitalizations

A total of 2952 individuals have been hospitalized due to COVID-19 in the county. **Figure 9** shows cumulative hospitalizations based on lab specimen collection date (as admission dates are incomplete). The highest number of hospitalizations continues to be among those ages 60-79 (N = 1094). The highest number of hospitalizations have occurred among the Black/AA community (N = 1144), followed by the Non-Hispanic White community (N = 1012) and then the Hispanic community (N = 564). Overall, counts are lower among other racial and ethnic groups. Females now slightly outnumber males, comprising 51.5% of all hospitalized cases. More individuals among lower income than higher income groups have been hospitalized, with a clear income gradient observed. Over the last week, we have seen an increase in hospitalizations among all income groups, those 60-79 and 80+, Non-Hispanic Black/AAs and Whites, and both sexes.

1600 1600 Median household income quartile Age 1400 1400 Q1: Lowest < 18 Q2: Low-medium 18-24 1200 1200 Q3: Medium-high 25-39 Q4: Highest 40-59 Cumulative cases among patients who became hospitalized 1000 1000 60-79 800 800 600 600 400 400 200 200 1600 1600 Race/Ethnicity* Gender 1400 1400 Black or AA Male White Female 1200 1200 Other/Unknown Hispanio . Asian 1000 1000 AIAN NHOPI 800 800 Multiple Races Other 600 600 400 400 200 200 \$\$\pa\fa\quad \text{\te}\text{\texi\texi{\text{\texi}\text{\text{\text{\text{\text{\text{\text{\text{\text{\texi}\texit{\text{

Figure 9: Cumulative hospitalizations in Milwaukee County

Date of specimen collection

Data source: Wisconsin Electronic Disease Surveillance System (WEDSS) Created by the Milwaukee County COVID-19 Epidemiology Intel Team


*Race and ethnicity were combined into one variable where the Hispanic category includes Hispanics of any race.

AIAN stands for American Indian or Alaska Native and NHOPI stands for Native Hawaiian or Other Pacific Islander.

When examined as population-based rates and case-based rates in **Figure 10**, hospitalization patterns are also apparent by demographic characteristics. Both population- and case-based hospitalization rates exhibit a clear age group gradient, with older age groups experiencing higher rates. By race and ethnicity, population and case-based hospitalization rates are highest among the Black/AA population and the population-based rate is lowest for non-Hispanic Whites. Rates by gender are very similar, with higher hospitalization rates among males. All rates presented are crude rates and only groups with 10 or more


total hospitalized cases are shown. Over the last week, we have seen a continued concerning uptick in the hospitalization rate among those 80+, and to a lesser degree, those 60-79.

Figure 10: Population and case based hospitalization rates in Milwaukee County




Age	N Hospitalized Cases	Rate per 1,000 residents	Rate per 100 cases
< 18	72	0.31	1.61
18-24	108	0.72	1.59
25-39	386	2.36	3.13
40-59	796	3.45	7.30
60-79	1094	7.66	22.22
80+	496	14.06	39.18

Specimen collection date



Race/Ethnicity*	N Hospitalized Cases	Rate per 1,000 residents	Rate per 100 cases
Black or AA	1144	4.59	12.20
White	1012	2.05	6.76
Hispanic	564	4.01	5.02
Asian	119	2.94	8.67
AIAN	20	4.30	10.53

Specimen collection date



Gender	N Hospitalized Cases	Rate per 1,000 residents	Rate per 100 cases
Male	1430	3.10	7.57
Female	1520	3.09	7.00

Specimen collection date

Data source: Wisconsin Electronic Disease Surveillance System (WEDSS) Created by the Milwaukee County COVID-19 Epidemiology Intel Team

*Race and ethnicity were combined into one variable where the Hispanic category includes Hispanics of any race.

AIAN stands for American Indian or Alaska Native and NHOPI stands for Native Hawaiian or Other Pacific Islander.

, in the standard of American Malari of American Standard of Stand

Deaths

There are now a total of 572 confirmed deaths in Milwaukee County, representing a case fatality rate of 1.4%. We observed 7 new deaths over the past week in the county. Mortality patterns differ by demographic characteristics, as shown in **Figure 11**. The largest number of deaths are recorded among those age 60 or older. The largest number of deaths are recorded for males (N = 293) and for non-Hispanic Whites (N = 271) followed by Black/AA residents (N = 203). By income, there are a larger number of deaths among the two lower income groups as compared to the two higher income groups. Deaths among Hispanics remain relatively low.

300 300 Median household income quartile Age Q1: Lowest < 18 250 250 Q2: Low-medium 18-24 Q3: Medium-high 25-39 200 Q4: Highest 200 40-59 60-79 150 150 100 100 50 50 0 7,76 71/16 Vidio, 1,10,15 AND!O 10, 10 10, 10 APTON Juno 'sep.on 'sepon Julio? nayay w in of 18 of 18 20, 40 0, 40 Cumulative deaths 300 300 Gender Race/Ethnicity* Black or AA 250 250 White Female Other/Unknown Hispanic 200 200 Asian AIAN NHOPI 150 150 Multiple Races Other 100 100 50 50

Figure 11: Cumulative deaths in Milwaukee County

Data source: Wisconsin Electronic Disease Surveillance System (WEDSS)
Created by the Milwaukee County COVID-19 Epidemiology Intel Team

to 0, 1,0 0, 1,0 0,1,0 0

*Race and ethnicity were combined into one variable where the Hispanic category includes Hispanics of any race.


AIAN stands for American Indian or Alaska Native and NHOPI stands for Native Hawaiian or Other Pacific Islander.

In terms of population- and case-based rates shown in **Figure 12**, there is a clear age category gradient, with higher death rates among older populations. Gender-based rates are very similar. Black/AA populations have the highest population and case-based death rates. All rates presented are crude rates and only groups with 10 or more total deaths are shown.


Date of death

to by the fleet mithing my my trafting to be on the by


Figure 12: Population and case based death rates in Milwaukee County



Age	N Deaths	Rate per 1,000 residents	Rate per 100 cases
18-39	10	0.03	0.05
40-59	66	0.29	0.61
60-79	250	1.75	5.08
80+	246	6.97	19.43



Race/Ethnicity*	N Deaths	Rate per 1,000 residents	Rate per 100 cases
Black or AA	203	0.82	2.16
White	271	0.55	1.81
Hispanic	71	0.51	0.63
Asian	17	0.42	1.24



Gender	N Deaths	Rate per 1,000 residents	Rate per 100 cases
Male	293	0.63	1.55
Female	278	0.56	1.28

Data source: Wisconsin Electronic Disease Surveillance System (WEDSS)

Created by the Milwaukee County COVID-19 Epidemiology Intel Team

*Race and ethnicity were combined into one variable where the Hispanic category includes Hispanics of any race.

AIAN stands for American Indian or Alaska Native and NHOPI stands for Native Hawaiian or Other Pacific Islander.

Testing Coverage

Testing for the novel coronavirus is an important public health response to limiting the spread of the infection. Testing capacity was limited in Milwaukee County and across the country earlier in the epidemic, but then increased. Since the first case of COVID-19 was diagnosed in Milwaukee County on March 6, 2020, a total of 553401 COVID-19 tests have been performed, with 506873 negative results and 46528 positive results. This represents a positive test rate of 8.4% since the beginning of the epidemic.

As shown in **Figure 13**, the total number of tests performed per week increased until early July and then peaked and declined, with an increase observed since early September. As shown in **Figure 14**, the percentage of positive tests has varied over the course of the epidemic, with a high of 25-30% in early April. The percentage of positive tests was 14.0% over the past week compared to 10.9% the previous week. **Figure 14** also illustrates the 14-day trend in the percent positive tests, showing a significant increase. Percent positive should be interpreted in the context of potential data delays given the large numbers of tests conducted in recent weeks, and considering that data entry for positive tests is prioritized.

35000 Number of tests performed Test result 30000 Negative Positive 25000 20000 15000 10000 5000 1,32° 1,3 20 21 July July July July 10 00

Figure 13: Milwaukee County number of tests per week

One-week window start date

Data source: Wisconsin Electronic Disease Surveillance System (WEDSS) Created by the Milwaukee County COVID-19 Epidemiology Intel Team




Figure 14: Milwaukee County percent positive tests

Date of specimen collection

Date of specimen collection


Data source: Wisconsin Electronic Disease Surveillance System (WEDSS)
Created by the Milwaukee County COVID-19 Epidemiology Intel Team

Spatial Patterns of Cases and Testing


COVID-19 spread is spatially patterned. **Map 1** below illustrates the cumulative burden (all confirmed cases) of COVID-19 in Milwaukee County. **Map 2** shows cases confirmed over the last week. **Map 3** shows the overall testing rate across the population. **Map 4** shows the testing rate over the last week. **Map 5** depicts the percentage of tests that were confirmed positive. **Map 6** shows cumulative COVID-19 related hospitalizations. **Map 7** shows the percentage of cases who have been hospitalized. All are crude rate maps created using census block group level COVID-19 data from WEDSS and population data from the US Census. The maps are smoothed to protect confidentiality and ensure that rates are stable while still providing geographic detail. Deciles are used to define categories. High rates are depicted in red with lower rates depicted in blue. Of note, some of the higher rates observed can be attributed to infections that have spread within group quarters, such as a nursing home, prison, or long-term care facility.

Decile Map 1: All confirmed cases of COVID-19

Created by the Milwaukee County Covid-19 Epidemiology Intel Team




Decile Map 2: Confirmed cases of COVID-19 within the last week




Method: A grid of points is used to estimate rates continuously across the map, based on the nearest cases with a minimum of 15 confirmed cases included.


Data Sources: Wisconsin Electronic Disease Surveillance System (WEDSS) (incidence data)

Decile Map 3: Overall testing rate



Decile Map 4: Testing rate within the last week






Method:
A grid of points is used to estimate rates continuously across the map, based on the nearest cases with a minimum of 15 tests included.


Data Sources: Wisconsin Electronic Disease Surveillance System (WEDSS) (incidence data) 2018 American Community Survey (population data) City of Milwaukee Map Milwaukee Portal (neighborhood boundaries) Census Bureau TIGER/Line Shapefiles (census place boundaries)

Created by the Milwaukee County Covid-19 Epidemiology Intel Team

Decile Map 5: Percentage of tests that were confirmed positive




Decile Map 6: COVID-19 related hospitalizations



Method:
A grid of points is used to estimate rates continuously across the map, based on the nearest cases with a minimum of 15 hospitalized cases included.

Data Sources: Wisconsin Electronic Disease Surveillance System (WEDSS) (incidence data)

Decile Map 7: Percentage of COVID-19 cases that were hospitalized



Method:
A grid of points is used to estimate rates continuously across the map, based on the nearest cases with a minimum of 15 confirmed cases included.

Data Sources: Wisconsin Electronic Disease Surveillance System (WEDSS) (incidence data)

Data Sources & Acknowledgments

This report was created by faculty and staff in the Medical College of Wisconsin (MCW) Institute for Health and Equity (IHE) in partnership with representatives from local health departments and faculty from the University of Wisconsin-Milwaukee Zilber School of Public Health. Data sources include the Wisconsin Electronic Disease Surveillance System (WEDSS), the US Census Bureau, the Milwaukee County Medical Examiner's office, the Emergency Medicine Resource, and publicly available data obtained from local health and emergency response agencies. Data from the Wisconsin Electronic Data Surveillance System (WEDSS) summarized for the week includes data from October 21, 2020 through October 27, 2020. This work was funded by the Advancing a Healthier Wisconsin Endowment at the Medical College of Wisconsin.

Contact Information

For additional questions on this report, please contact Darren Rausch, Health Officer/Director, Greenfield Health Department, and Lead, Milwaukee County COVID-19 Epidemiology Intel Team: Darren.Rausch@greenfieldwi.us or (414) 329-5275.